


The astrophysics of extreme mass-ratio inspiral sources

Clovis Hopman
Weizmann, Leiden



Overview

- ✓ Why EMRIs?
- ✓ Stellar dynamics and content near massive black holes
- ✓ Inspiral dynamics, loss-cone theory
- ✓ Dynamics within 0.01 pc: mass-segregation and resonant relaxation
- ✓ Alternative routes to EMRIs

Why are EMRIs important?

- ✓ Most detailed tests for general relativity
- ✓ Stellar content very close to massive black holes
- ✓ Stellar dynamics very near massive black holes
- ✓ Mass distribution of massive black hole
- ✓ Spin distribution of massive black holes; merger history
- ✓ Accretion disks
- ✓ Do intermediate mass black holes exist?
- ✓ Are massive black holes really black holes? (Or, let's say, boson stars?)

Stars near massive black holes




- ✓ Very dense environment of stars
- ✓ Density diverges as a power-law: $n \sim r^{-(1.5-2)}$
- ✓ Short relaxation times ($t_r < 10$ Gyr)
- ✓ Black hole dominates dynamics within ~ 1 pc.

Infra-red observations of the Galactic center


Bahcall & Wolf 1976, 1977; Alexander 1999; Genzel et al. 2003

Stars near massive black holes




Infra-red observations of the Galactic center

- ✓ Very dense environment of stars
- ✓ Density diverges as a power-law: $n \sim r^{-(1.5-2)}$
- ✓ Short relaxation times ($t_r < 10$ Gyr)
- ✓ Black hole dominates dynamics within ~ 1 pc.




Bahcall & Wolf 1976, 1977; Alexander 1999; Genzel et al. 2003

How to get near a massive BH



Hills & Bender;
Sigurdsson & Rees;
Miralda-Escude &
Gould; Gair; Freitag;
Ivanov; Hopman &


How to get near a massive BH



Bursts of GWs
may be
observable in our
own Galactic
center (Rubbo et
al. 2006)


Hills & Bender;
Sigurdsson & Rees;
Miralda-Escude &
Gould; Gair; Freitag;
Ivanov; Hopman &

Loss-cone theory




Lightman & Shapiro (1977)

Loss-cone theory



Kick regime:
$$\Gamma \sim \frac{N\vartheta^2}{P}$$


Loss-cone theory



Kick regime: $\Gamma \sim \frac{N\vartheta^2}{P}$

Diffusion regime: $\Gamma \sim \frac{N}{t_r}$


Angular momentum scattering and energy dissipation



Hils & Bender (1995); Hopman & Alexander (2005)

Angular momentum scattering and energy dissipation


Critical scale determined by equating inspiral time and scatter time near the LSO



Hils & Bender (1995); Hopman & Alexander (2005)

EMRI event rate



- ✓ Critical distance at $a_{\text{GW}} = 0.01$ pc
- ✓ Nature of EMRIs determined by dynamics $< a_{\text{GW}}$
- ✓ Event rate: integrate diffusion expression over inspiral probability function:



Hopman & Alexander (2005)

$$\Gamma_i = f_s \int_0^\infty \frac{da N(a) S(a)}{\ln(J_m/J_{lc}) t_r(a)}$$

Dynamics within a_{GW} : mass segregation





Stellar BHs sink to center and dominate dynamics within a_{GW} .
Relaxation time decreases due to presence of stellar BHs.

Hopman & Alexander (2006)

Freitag, Amaro-Seoane & Kalogera (2006)

Dynamics within a_{GW} : resonant relaxation




Resonant relaxation increases EMRI rate by factor
Crucial to include general relativity!

Rauch & Tremaine (1995)
Rauch & Ingalls (1998)


Hopman & Alexander (2000)

LISA EMRIs: high eccentricities at $P=3$ hrs

Massive Black Holes:




Intermediate Mass Black Holes:



Hopman & Alexander (2005)


Orbits must be modeled (pseudo-) relativistically
Hopman & Alexander 2005; Gair et al. 2005

Indirect EMRI capture mechanisms



Stars can be formed in accretion disks and spiral in due to interaction with gas (Levin 2003, 2006)


Indirect EMRI capture mechanisms



Stars can be formed in accretion disks and spiral in due to interaction with gas (Levin 2003, 2006)

Binary stars are tidally disrupted, forming a hyper-velocity star and an EMRI (Miller, Freitag & Hamilton 2005)

Indirect EMRI capture mechanisms



Stars can be formed in accretion disks and spiral in due to interaction with gas (Levin 2003, 2006)

Binary stars are tidally disrupted, forming a hyper-velocity star and an EMRI (Miller, Freitag & Hamilton 2005)

Efficient loss-cone refilling by triaxiality (Holley-Bockelmann et al. 2006) and massive perturbers (Perets, Hopman & Alexander 2006)

Indirect EMRI capture mechanisms




Stars can be formed in accretion disks and spiral in due to interaction with gas (Levin 2003, 2006)

Binary stars are tidally disrupted, forming a hyper-velocity star and an EMRI (Miller, Freitag & Hamilton 2005)

Stars can be tidally captured by intermediate mass BHs. After an ultraluminous X-ray phase, they leave the main sequence and spiral in (Holley-Bockelmann et al. 2006) and massive perturbers (Perets, Hopman & Alexander 2006) (Hopman et al 2004, Hopman & Portegies Zwart 2005)

A bimodal distribution of eccentricities?



Conclusions

- ✓ Direct captures: event rate of $\sim 100/\text{Gyr}/\text{MBH}$, but rather uncertain. Stellar BHs dominate rate
- ✓ Rates depend on dynamics within $a_{\text{GW}} \sim 0.01$ pc: mass-segregation, resonant relaxation, ...
- ✓ Direct capture sources are eccentric
- ✓ Indirect capture leads to low eccentricity EMRIs